

レジメン名	商品名	剤形	投与量	スケジュール (日)																											
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
FP	フルオロウラシル	注	800 mg/m ²	○	○	○	○	○	→																						
	シスプラチン	注	80 mg/m ²	○																											
	備考	フルオロウラシル：120時間持続投与																													
	副作用	骨髄抑制、吐き気、味覚障害、腎機能障害など																													
5-FU+アクプラ +RT	フルオロウラシル	注	800 mg/m ²	○	○	○	○	○	→																						
	アクプラ	注	80 mg/m ²	○																											
	備考	フルオロウラシル：96時間持続投与																													
	副作用	骨髄抑制、吐き気、味覚障害、腎機能障害など																													
オプジーボ (食道、4週毎)	オプジーボ	注	240 mg	○																											
	備考																														
	副作用	免疫関連有害事象																													
FP+RT (50.4Gy)	フルオロウラシル	注	1000 mg/m ²	○	○	○	○	○	→																						
	シスプラチン	注	75 mg/m ²	○																											
	備考	フルオロウラシル：96時間持続投与																													
	副作用	骨髄抑制、吐き気、味覚障害、腎機能障害など																													

投与間隔 : 49 日

レジメン名	商品名	剤形	投与量	スケジュール (日)																											
				1	2	...	7	8	9	...	14	15	16	...	21	22	23	...	28	29	30	...	35	36	37	49
PTX Weekly (B法)	パクリタキセル	注	100 mg/m ²	○						○																					
	備考																														
	副作用	骨髄抑制、脱毛、しびれ、アレルギー、関節痛・筋肉痛など																													